

Structure of fermion nodes and nodal cells

Lubos Mitas¹

¹*Department of Physics and CHiPS, North Carolina State University, Raleigh, NC 27695*

(Dated: February 6, 2008)

We study nodes of fermionic ground state wave functions. For $2D$ and higher we prove that spin-polarized, noninteracting fermions in a harmonic well have two nodal cells for arbitrary system size. The result extends to other noninteracting/mean-field models such as fermions on a sphere, in a periodic box or in Hartree-Fock atomic states. Spin-unpolarized noninteracting states have multiple nodal cells, however, interactions and many-body correlations generally relax the multiple cells to the minimal number of two. With some conditions, this is proved for interacting $2D$ and higher dimensions harmonic fermion systems of arbitrary size using the Bardeen-Cooper-Schrieffer variational wave function. Implications and extent of these results are briefly discussed.

PACS numbers: 02.70.Ss, 03.65.Ge

A fermion node is a subspace of fermion configurations for which a real wave function describing the fermionic system vanishes due to the antisymmetry. In general, for N spin-polarized fermions in d dimensions the fermion node is a $(dN - 1)$ -dimensional hypersurface given by an implicit equation $\Psi(R) = 0$, where Ψ is the wave function with fermion coordinates $R = (\mathbf{r}_1, \dots, \mathbf{r}_N)$. The location of the nodal manifold is of key importance for quantum Monte Carlo (QMC) methods [1, 2, 3, 4] since the exact node enables us to solve the stationary Schrödinger equation with computer time scaling as a low-order polynomial in N . Remarkably, even rather crude Hartree-Fock (HF) or post-HF wave function nodes, routinely used in the fixed-node approximation QMC methods, provide $\approx 95\%$ of the correlation energy in a variety of systems with hundreds of valence electrons [1, 2, 3, 4]. However, to reach beyond this level of accuracy has proved to be challenging because of limited understanding of fermion nodes [4, 5, 6, 7, 8, 9].

The exact nodes for interacting systems are known only for a very few two-electron triplet atomic states [5, 10] and, very recently, the exact node of three-electron $^4S(p^3)$ state has been discovered [11]. It turned out that the nodes of these high-symmetry states have rather simple topologies and divide the configuration space into two compact nodal cells in which the wave function is positive or negative ("plus" or "minus" cell). The two nodal cells were found also in noninteracting spin-polarized $2D$ and $3D$ fermions with up to 200 particles using a numerical proof [5]. The same noninteracting but spin-unpolarized systems trivially have four nodal cells since the wave function is a product of spin-up and spin-down determinants. Interestingly, analysis of interacting unpolarized few-particle systems [6, 9, 12] has revealed that the electron correlation can change the node topologies and number of nodal cells. For example, the correlated wave functions of the Be atom or N_2 molecule exhibit two nodal cells while the corresponding Hartree-Fock ones have four. It was therefore conjectured that the bisection of the configuration space into the two nodal cells might

be a generic property of fermionic ground states (with exceptions discussed later). In this Letter we prove that for $2D$ spin-polarized noninteracting harmonic fermions the ground state node divides the configuration space into the minimal number of two nodal cells for any system size. The proof method extends to higher dimensions and carries over to other models such as fermions on a sphere, in a periodic box, for atomic states, etc. We show that the same holds, in general, for interacting systems using correlated Bardeen-Cooper-Schrieffer wave functions and we briefly discuss the implications.

We recall two basic properties of fermion nodes derived by Ceperley [5]. a) Nondegenerate ground state wave functions satisfy the so-called *tiling property* which states that by applying all possible particle permutations to an arbitrary nodal cell one covers the entire configuration space. Note that this does not specify the number of nodal cells. b) Consider three particles i, j, k in a spin-polarized system with wave function $\Psi(R)$. We call the particles i, j, k connected, if there exists a triple exchange $ijk \rightarrow jki$ path that does not cross the node, ie, $|\Psi(R)| > 0$ along the exchange path. More connected particles, such as the following six ones, can form a cluster: . If there exists a point R_t such that triple exchanges *connect all the particles into a single cluster* then $\Psi(R)$ has only *two nodal cells*. The tiling property implies that once the particles are connected for R_t the same applies to any point in the cell as further explained in Ref. [5].

We illustrate the properties of nodes in $1D$ using spin-polarized fermions in a $1D$ harmonic oscillator well. The wave function is a Slater determinant $\Psi(1, \dots, N) = \det\{\phi_k(\mathbf{r}_i)\} = A \prod_i e^{-x_i^2/2} \det\{1, 2x, \dots, H_{N-1}(x)\}$ where $H_n(x)$ is a Hermite polynomial of degree n and A is the normalization. We omit the prefactors and transform the Slater matrix to monomials so that the wave function is given by the Vandermonde determinant

$$\Psi_{1D}(1, \dots, N) = \det\{1, x, x^2, \dots, x^{N-1}\} = \prod_{i < j} (x_j - x_i) \quad (1)$$

The node is encountered whenever two fermions pass through each other and the wave function has $N!$ nodal cells since any permutation requires at least one node-crossing. In general, the derived node is exact for other $1D$ models including systems with interactions.

Now consider spin-polarized fermions in a $2D$ harmonic well. The one-particle states are simply $\phi_{nm} = C_{nm}H_n(x)H_m(y)$, $n, m = 0, 1, \dots$ where C_{nm} includes the gaussian and normalization which are absorbed into a common prefactor and omitted. The Slater matrix elements can be rearranged to monomials and we write

$$\Psi_{2D}(1, \dots, N) = \det\{1, x, y, \dots, x^n y^m, \dots\} \quad (2)$$

The closed-shell states and the system size are labeled by $M = 1, 2, \dots$ where $n + m \leq M$ with the number of fermions given by $N = (M + 1)(M + 2)/2$.

Using induction we prove that the wave function in Eq. 2 has only two nodal cells for any $M > 0$. This is indeed true for $M = 1$ with three-particle wave function $\Psi_{2D}(1, 2, 3) = \det\{1, x, y\}$. In order to show this, it is convenient to extend the particle $2D$ coordinates by a "dummy" third dimension as $\mathbf{r}_i = (x_i, y_i, 0)$. Then $\Psi_{2D}(1, 2, 3) = \mathbf{z}_0 \cdot (\mathbf{r}_{21} \times \mathbf{r}_{31})$ where $\mathbf{z}_0$ is the unit vector in the third dimension and $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$. Clearly, there are only two nodal cells since the set of vectors $\mathbf{z}_0, \mathbf{r}_{21}, \mathbf{r}_{31}$ is either left- or right-handed, and the node is encountered whenever the three particles are collinear, ie, $\mathbf{r}_{21} \times \mathbf{r}_{31} = 0$. Also, the particles are connected by triple exchanges without node-crossing (eg, rotate an equilateral triangle).




FIG. 1: Particles arranged in a Pascal-like triangle pattern. The circles denote the particle positions and inside the circles are particle labels.

We now consider a general system with $M > 1$ and arrange the particles into a Pascal-like triangle pattern on a rectangular mesh as shown in Fig.1. For this arrangement the determinant can be explicitly evaluated for any M by subsequent factorization of lines of particles [13]. Due to the space constraints and to simplify the notation we illustrate the factorization on a few-particle example; the generalization to an arbitrary size is straightforward.

For example, for $M = 2$ the wave function reads

$$\Psi_{2D}(1, \dots, 6) = \det\{1, x, y, x^2, xy, y^2\} = \begin{vmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ x_0 & x_1 & x_0 & x_2 & x_1 & x_0 \\ y_0 & y_0 & y_1 & y_0 & y_1 & y_2 \\ x_0^2 & x_1^2 & x_0^2 & x_2^2 & x_1^2 & x_0^2 \\ x_0 y_0 & x_1 y_0 & x_0 y_1 & x_2 y_0 & x_1 y_1 & x_0 y_2 \\ y_0^2 & y_0^2 & y_1^2 & y_0^2 & y_1^2 & y_2^2 \end{vmatrix} \quad (3)$$

where the particle positions are given in Fig.1 (our point R_t). Clearly, all the last-row elements containing y_0 can be eliminated by adding a multiple of the third row. In a similar way we eliminate all the matrix elements containing y_0 , and the determinant factorizes as

$$\Psi_{2D}(1, \dots, 6) = \prod_{i=1}^2 (y_i - y_0)^{n_i} \Psi_{1D}(1, 2, 4) \Psi_{2D}(3, 5, 6) \quad (4)$$

where n_i is the number of particles on line $y = y_i$. Note that one of the factors is the $2D$ wave function for the system with M reduced by one. The same factorization structure is obtained for an arbitrary M and, obviously, it can be applied recursively. The wave function for a general size M is then given by

$$\begin{aligned} \Psi_{2D}(1, \dots, N) &= \pm \prod_{i>0}^M (y_i - y_0)^{n_i} \Psi_{1D}(I_0) \Psi_{2D}(1, \dots, N/I_0) \\ &= \pm \prod_{l=0}^{M-1} \left[\prod_{i>l}^M (y_i - y_l)^{n_i} \Psi_{1D}(I_l) \right] \end{aligned} \quad (5)$$

where $I_l = i_1^{(l)}, \dots, i_{M+1-l}^{(l)}$ denotes the labels of particles lying on the line l while $(1, \dots, N/I_0)$ means that the labels in I_0 are omitted from $1, \dots, N$. The sign depends on the number of row exchanges and on the actual ordering of particles. Note the translation invariance of Eq. 5.

For the induction step we assume that triple exchanges connect all the particles in the system of size M . Let us show that the same is true for the system of size $M + 1$ (see Fig.1). We first factorize out the line $y = y_0$ as given by Eq. 5. One of the factors is the $2D$ wave function for the system of size M , which contains all additional particles except the one with coordinates $[x_{M+1}, y_0]$. By the original assumption, such a system has all the particles connected. If instead of the horizontal line we factorize out the vertical line ($x = x_0$) we see that the particle at $[x_{M+1}, y_0]$ is also connected, thus concluding the proof.

The proof can be generalized in several ways. The same arguments apply to $3D$ since particles can be arranged into an appropriate $3D$ pattern and the wave function evaluated by recursive factorization of *planes and lines*.

In fact, the result holds and the two nodal cell property is correct for arbitrary dimension $d > 1$! The proof applies also to other noninteracting or mean-field models with polynomial entries in the Slater matrix such as fermions on a sphere or in a periodic box with details given elsewhere [14]. The proof can be modified at least for some open-shell states as well while taking into account node ambiguities if there are degeneracies [5, 8].

The proof method can be further combined with symmetries for cases when a complete factorization is not obvious, such as for multi-shell atomic states. For example, spin-polarized ground states $1s2s2p^n$, $n = 1 - 3$ (and beyond) also have two nodal regions. We show this for the ${}^6S(1s2s3p^3)$ state in the noninteracting and HF limits. The wave function is written as $\Psi_{at}(1, \dots, 5) = \det\{\rho_{1s}^*(r), \rho_{2s}^*(r), x, y, z\}$ where $\rho_{1s}^*(r) = \rho_{1s}(r)/\rho_{2p}(r)$ and $\rho_{2s}^*(r) = \rho_{2s}(r)/\rho_{2p}(r)$ since nonnegative $\rho_{2p}(r)$ is factorized out. The dimensionless coordinates are rescaled by the atomic number Z and the Bohr radius a_0 as $\mathbf{r} \leftarrow Z\mathbf{r}/a_0$. Let us place particle 1 at the origin and particles 2-5 on the surface of a sphere with the radius η_0 equal to the radial node of $\rho_{2s}(r)$ orbital, ie, $\rho_{2s}(\eta_0) = 0$. For such configurations we obtain

$$\Psi_{at}(1, \dots, 5) = \rho_{1s}^*(\eta_0)\rho_{2s}^*(0)\mathbf{r}_{32} \cdot (\mathbf{r}_{42} \times \mathbf{r}_{52})$$

so that any three-particle exchange from 2,3,4,5 easily avoids node-crossing by appropriate positioning and rotations on the sphere. The particle 1 is connected by the exchange $123 \rightarrow 312$ parametrized as $\mathbf{r}_1(t) = \eta_0[t, 0, 0]$; $\mathbf{r}_2(t) = \eta_0[c(t), s(t), 0]$ and $\mathbf{r}_3(t) = \eta_0[0, 1-t, 0]$ where $t = 0$ ($t = 1$) corresponds to the beginning (end) point of the exchange path while $c(t) = \cos(\pi t/2)$ and $s(t) = \sin(\pi t/2)$. Setting $\mathbf{r}_4 = [0, 0, \eta_0]$ and $\mathbf{r}_5 = [0, 0, -\eta_0]$ we find that Ψ_{at} is proportional to

$$\rho_{2s}^*(t\eta_0)c(t)(1-t) + \rho_{2s}^*[(1-t)\eta_0]s(t)t > 0$$

The inequality holds for the whole path $0 \leq t \leq 1$ since $\rho_{2s}^*(t\eta_0) > 0$ for $0 \leq t < 1$ for both noninteracting and HF cases. The proof can be further extended to more shells such as $1s2s2p^33s3p^3$ and $1s2s2p^33s3p^33d^5$ [14].

Spin-unpolarized systems. Due to the product of spin-up and -down determinants, the number of nodal cells in noninteracting unpolarized systems is twice the number of cells of the half-filled spin-polarized counterparts. The proof for atomic states above then implies that the HF wave functions for atoms with $Z > 3$ up to the third-row elements have four nodal cells.

Interactions in spin-polarized systems. In general, the shape and topology of the nodal manifold is influenced by interactions and many-body effects. Consider the lowest atomic quartet of S symmetry and even parity ${}^4S^e(1s2s3s)$ (clearly not the lowest quartet, which is odd ${}^4P^o(1s2s2p)$). The noninteracting wave function is given by $\Psi_{at}(1, 2, 3) = \det\{\rho_{1s}(r), \rho_{2s}(r), \rho_{3s}(r)\}$ and has *six* nodal cells. Since $\Psi_{at}(1, 2, 3)$ depends only on

distances it is quasi-1D and the nodes are the same as the ones given by Eq. 1. For the interacting system the correlation is included by adding to the wave function the lowest (and dominant) double excitation $2s3s \rightarrow 2p_x3p_x + 2p_y3p_y + 2p_z3p_z$ with a weight w . The correlated wave function then allows for exchanges without node-crossing. Define $\mathbf{r}_a(t) = [0, c(t), s(t)]$, $\mathbf{r}_b(t) = [c(t), s(t), 0]$, $\mathbf{r}_{ab}(t) = [1 - g(t)]\mathbf{r}_b(t) - g(t)\mathbf{r}_a(t)$, $\mathbf{r}_c(t) = \mathbf{r}_{ab}(t)/|\mathbf{r}_{ab}(t)|$, and $\mathbf{r}_d(t) = \mathbf{r}_c(t) \times \mathbf{r}_a(t)$, where $g(t) = 3t(1-t)$. The exchange path $123 \rightarrow 231$ is then $\mathbf{r}_3(t) = r_3(t)\mathbf{r}_a(t)$, $\mathbf{r}_2(t) = r_2(t)\mathbf{r}_c(t)$ and $\mathbf{r}_1(t) = r_1(t)\mathbf{r}_d(t)/|\mathbf{r}_d(t)|$. The radial parts are given by $r_1(t) = \eta_m + q(2t-1)$, $r_2(t) = \eta_m + q(1-t)$ and $r_3(t) = \eta_m - qt$. η_m is the mean value of the radial node of ρ_{2s} and the first radial node of ρ_{3s} orbitals while $0 < q < a_0|w|$ (for the Coulomb e-e interaction w is ≈ -0.05). The path is orchestrated so that in the region where the noninteracting component vanishes the correlation dominates and the particles become connected. This illustrates two points: imposing symmetries at the mean-field level can generate multiple cells and, in general, for $d > 1$ the interactions lift this "nodal cell degeneracy" and relax multiple cells to the minimal two.

Interactions in spin-unpolarized systems. The change from four to two nodal cells due to interactions has been demonstrated for the first time on the Be atom [9] using the two-configuration correlated wave function and the connected cluster construction adapted to spin-unpolarized systems. Consider a simultaneous exchange of one (or more) pair(s) of spin-up particles and one (or more) pair(s) of spin-down particles. For noninteracting wave functions such simultaneous pair exchanges imply that the node will be crossed once or multiple times. If there exists a point R_f such that during the simultaneous spin-up and -down pair exchanges the inequality $|\Psi| > 0$ holds along the whole path, then the wave function has only two nodal cells.

Consider a singlet state of $2N$ particles in a 2D harmonic well with particles interacting by pair potential. With some restrictions, we show that the correlation included in the Bardeen-Cooper-Schrieffer (BCS) pairing wave function [7, 15] given by $\Psi_{BCS}(1, \dots, 2N) = \det[\Phi(i, j)]$ is enough to eliminate the noninteracting four nodal cells and fuse them into the minimal two. Here $\Phi(i, j) = \Phi(j, i)$ is a singlet pair orbital for $i \uparrow$ and $j \downarrow$ fermions and we decompose it into noninteracting and correlated components $\Phi(i, j) = \Phi_0(i, j) + \Phi_{corr}(i, j)$. Using one-particle orbitals we can write $\Phi_0(i, j) = \sum_{n+m=0}^M \phi_{nm}(i)\phi_{nm}(j)$ while $\Phi_{corr}(i, j) = \sum_{n+m>M} c_{nm}\phi_{nm}(i)\phi_{nm}(j)$ where $\{c_{nm}\}$ are variational parameters.

We illustrate this first on a six-particle singlet, $M = 1$, $2N = 6$. In cylindric coordinates (r, φ) we find $\Phi_0(i, j) = 1 + 2r_i r_j \cos \varphi_{ij}$ where $\varphi_{ij} = \varphi_i - \varphi_j$, with the gaussians and normalization omitted. $\Phi_{corr}(i, j)$ is constructed as a sum of orbitals from the next unoccupied shell ($n+m =$




FIG. 2: (a) Positions of six 2D harmonic fermions with two particles at the origin and two pairs on circles with radii r_a, r_b . (b) Positions of the spin-up particles for the $M = 2, 2N = 12$ particle singlet. The spin-down positions pattern is similar except for the different radii and angles.

2) and we find $\Phi_{corr}(i, j) = \alpha[2(r_i r_j \cos \varphi_{ij})^2 - r_i^2 - r_j^2]$, where a variational parameter α has been included. We place the particles as in Fig.2(a). For such a configuration $\Psi_{BCS}(1, \dots, 6) = 8\alpha r_a r_b \cos \varphi [2(r_a r_b \cos \varphi)^2 - r_a^2 - r_b^2]$. The rotation of the system by π exchanges the pair of particles for each spin, and since the wave function is rotationally invariant, it is enough to show that it is nonzero for a single point, eg, $r_a = r_b = 1, \varphi = \pi/4$. For $\alpha = 0$ the wave function vanishes since the particles lie on the noninteracting node. However, for any interaction which does not break the rotation invariance and gives $\alpha \neq 0$, the BCS wave function has only two nodal cells.

Remarkably, this can be generalized to an arbitrary size. Assume a closed-shell singlet with the total number of particles $2N = (M + 1)(M + 2)$ where N is even (for N odd the derivations are the same after placing one particle of each spin to the origin, see Fig.2(a)). We form $N/2$ pairs of particles in each spin subspace so that, say, the pair $i \uparrow, (i + 1) \uparrow$ has coordinates given by $(r_i, \varphi_i) = (r_k, \varphi_k)$, $(r_{i+1}, \varphi_{i+1}) = (r_k, \varphi_k + \pi)$, see Fig. 2(b). Here $k = 1, \dots, N/2$ labels the pairs in the spin-up channel; the spin-down particles are placed similarly and labeled by the pair index $l = N/2 + 1, \dots, N$. In this configuration the particles lie on the noninteracting node since $\det\{\Phi_0(i, j)\} = \det\{\sum_{n+m \leq M} \phi_{nm}(i)\phi_{nm}(j)\} = \det\{\phi_{nm}(i)\}\det\{\phi_{nm}(j)\}$ and the rotation of the system by π crosses the nodes in both spin channels so that both Slater determinants vanish due to the rotation invariance. Now, if all the N pair distances and angles $r_k, r_l, \varphi_k, \varphi_l$ are distinct, then each of the matrices $\{\phi_{nm}(i)\}, \{\phi_{nm}(j)\}$ has exactly one linearly dependent row, ie, their ranks are $N - 1$. This can be verified directly for small values of M and then using induction for any M . Consequently, the matrix $\{\Phi_0(i, j)\}$ has linear dependence in one row and one column, ie, it has the rank of $N - 1$ as well. In general, adding virtual states through $\Phi_{corr}(i, j)$ provides independent rows/columns

(eg, $M + 1$ independent rows/columns from the first unoccupied shell) which eliminate linear dependency so that $\det\{\Phi_0(i, j) + \Phi_{corr}(i, j)\}$ is nonzero. Assuming that the interactions do not break the rotation invariance, the correlated BCS wave functions have only two nodal cells. In fact, this can hold even if the invariance is broken but one would need to show it for the entire exchange path, not only for a single point. The proof extends to $d > 2$ and to other models as well, with details given elsewhere [14].

For the classes of fermion systems studied in this work the two nodal cell property indeed appears as a generic feature. That brings up an interesting question: when might this property not apply? We mention just some of the possibilities: i) additional symmetries and/or boundary conditions can generate additional nodal cells; ii) nonlocal or very strong/singular interactions can reorder the states (eg, an excited state becoming the ground state) or significantly change the nodes; iii) for systems with strong correlations, large degeneracies at the Fermi level, or at quantum phase transitions the properties of nodes are far from clear and require further study.

We believe the presented analysis and proofs provide a significant step forward in our understanding of topological properties of fermionic wave functions. The results offer an elegant and unifying framework for several previously conjectured or numerically investigated features and apply to arbitrary size for both noninteracting and interacting systems. The presented techniques extend to other models and open exciting perspectives for studies of many-body effects which are currently out of reach.

I would like to acknowledge the support by NSF DMR-0102668, DMR-0121361 and EAR-0530110 grants.

-
- [1] D.M. Ceperley and M.H. Kalos, in *Monte Carlo Methods in Statistical Physics*, Ed. K. Binder, pp.145-194, Springer, Berlin, 1979; K.E. Schmidt and D. M. Ceperley, in *Monte Carlo Methods in Statistical Physics II*, pp.279-355, Ed. K. Binder, Springer, Berlin, 1984.
 - [2] B.L. Hammond, W.A. Lester, Jr., and P.J. Reynolds, *Monte Carlo Methods in ab initio quantum chemistry*, World Scientific, Singapore, 1994.
 - [3] M.W.C. Foulkes, L. Mitás, R.J. Needs and G. Rajagopal, *Rev. Mod. Phys.* **73**, pp. 33-83 (2001).
 - [4] J. B. Anderson, *J. Chem. Phys.* **65** 4121 (1976); J. B. Anderson, *Phys. Rev. A*, **35**, 3550 (1987).
 - [5] D. M. Ceperley, *J. Stat. Phys.* **63**, 1237, (1991).
 - [6] W. A. Glauser, W. R. Brown, W. A. Lester, D. Bressanini, B. L. Hammond and M. L. Koszykowski, *J. Chem. Phys.* **97**, 9200 (1992).
 - [7] M. Casula and S. Sorella, *J. Chem. Phys.* **119**, 6500 (2003)
 - [8] W. M. C. Foulkes, Randolph Q. Hood and R. J. Needs, *Phys. Rev. B* **60**, 4558 (1999).
 - [9] D. Bressanini, D. M. Ceperley and P. Reynolds, in *Recent Advances in Quantum Monte Carlo Methods, II* Ed.

- W. A. Lester, S. M. Rothstein, and S. Tanaka, World Scientific, Singapore (2002).
- [10] D. Bressanini, and P. Reynolds, Phys. Rev. Lett. **95**, 110201 (2005).
 - [11] M. Bajdich, L. Mitas, G. Drobny, and L. K. Wagner, Phys. Rev. B **72**, 075131 (2005)
 - [12] L. Mitas, G. Drobny, M. Bajdich, L.K. Wagner, in *Condensed Matter Theories, Vol.20*, Ed. J.W. Clark, 2005, in press.
 - [13] C. K. Chui and L. M. Lai, in *Nonlinear and Convex Analysis*, Eds. B.L.Lin and S.Simons, pp. 23-35, Marcel Dekker, 1987.
 - [14] L. Mitas, to be published.
 - [15] J. Carlson, S.-Y. Chang, V. R. Pandharipande, and K. E. Schmidt, Phys. Rev. Lett. **91**, 050401 (2003)